

# Case Report Form CovidSurg-Cancer - Pancreas

NB: Additional data points may be required for specific cancer types



NIHR Global Health Research Unit on  
Global Surgery



Patient REDCap ID: \_\_\_\_\_

Age: 0-4w | 4-52w | 1-9y | 10-16y | 17-19y | 20-29y | 30-39y | 40-49y | 50-59y | 60-69y | 70-79y | 80-89y | 90y+ Sex: Female | Male ASA Grade: 1 | 2 | 3 | 4 | 5. Weight (<52 weeks only): \_\_\_\_\_ kg

WHO/ECOG Performance status: 0 | 1 | 2 | 3 | 4

BMI:

- |  |   |
|--|---|
| <input type="checkbox"/> Underweight (<18.5)               | <input type="checkbox"/> Moderately obese (30-34.9) |
| <input type="checkbox"/> Normal/healthy weight (18.5-24.9) | <input type="checkbox"/> Severely obese (35-39.9)   |
| <input type="checkbox"/> Overweight (25-29.9)              | <input type="checkbox"/> Very severely obese (≥40)  |

Comorbidities:

- |   |   |
|---|---|
| <input type="checkbox"/> Current smoker                           | <input type="checkbox"/> Congestive heart failure |
| <input type="checkbox"/> Asthma                                   | <input type="checkbox"/> Dementia                 |
| <input type="checkbox"/> Chronic Kidney Disease (Moderate/Severe) | <input type="checkbox"/> Diabetes Mellitus        |
| <input type="checkbox"/> COPD                                     | <input type="checkbox"/> Hypertension             |
| <input type="checkbox"/> Congenital abn (cardiac)                 | <input type="checkbox"/> Myocardial Infarction    |
| <input type="checkbox"/> Congenital abn (non cardiac)             | <input type="checkbox"/> Peripheral Vascular Dis  |
|   | <input type="checkbox"/> Stroke/TIA               |
|   | <input type="checkbox"/> Other: _____             |

Cancer-specific details:

**NCCN classification:**

Resectable | Borderline resectable (vein involvement) | Borderline resectable (arterial involvement) | Locally advanced

Date of cancer diagnosis: \_\_\_\_/\_\_\_\_/\_\_\_\_

Date of decision for curative surgery: \_\_\_\_/\_\_\_\_/\_\_\_\_

**Prior to COVID-19, did your hospital routinely operate on jaundiced patients with resectable PDAC without preoperative biliary drainage?**

- ☐ Yes ☐ No

**Was this patient potentially eligible for surgery without preoperative biliary drainage (PBD) at your unit?**

- ☐ Yes ☐ No

**Did the patient undergo PDB?**

- ☐ Yes – due to COVID-19 pandemic  
☐ Yes – unrelated to COVID-19 pandemic  
☐ No – jaundiced patient underwent surgery without PBD  
☐ No – patient not jaundiced

**Was the initial MDT (tumour board) decision for primary surgical treatment?**

Yes – decision for surgical Rx (optimal treatment option) | Yes – decision for surgical Rx (compromised option due to COVID-19) | No – decision for non-surgical Rx (optimal treatment option) | No – decision for non-surgical Rx (compromised option due to COVID-19)

**Reason of compromise:** \_\_\_\_\_

**Did the patient have an operation related to this cancer during the 3-month study window?** No / Yes

**If a cancer operation WAS performed:**

Date of surgery: \_\_\_\_/\_\_\_\_/\_\_\_\_

**Op performed in:** Dedicated COVID-free hospital | Dedicated COVID Rx hospital | Undesignated hospital type with ED | Undesignated hospital type without ED | Other: \_\_\_\_\_

**COVID-19 level:** 0 | 1 | 2 | 3 | 4 | 5

Did the patient have a **\*resolved\* COVID-19 infection** before the time of surgery? Yes – lab test/CT thorax | Clinical suspicion | No

Was **COVID-19 screening** performed preoperatively?

Laboratory test | CT thorax | Other: \_\_\_\_\_

Did patient have mandatory self-isolation before surgery?

Yes, > 2 weeks | Yes - < 2 weeks | No

**COVID-19 suspected at time of surgery?** Yes | No

**Tests performed to investigate SARS-CoV-2 status:**

	CT	CT	Swab	Swab
	(neg)	(pos)	(neg)	(pos)

4-7 days prior surgery ☐ ☐ ☐ ☐

1-3 days prior surgery ☐ ☐ ☐ ☐

Immediate | Urgent | Expedited | Elective

**If a cancer operation WAS performed:**

If **emergency cancer surgery** was required, why?

☐ Gastro-intestinal obstruction ☐ Bleeding ☐ Sepsis

☐ Tumour progression ☐ Organ perforation

☐ Other: \_\_\_\_\_

**Operation performed?** \_\_\_\_\_

**Approach:** Open | Minimally invasive

**Surgical intent:** Curative | Palliative

**Resection margin status:** R0 | R1 | R2 | Unknown

**Did this represent a change to your typical operative approach in the pre-COVID-19 era?**

- ☐ No change to operative approach  
☐ Yes, chose to avoid minimally invasive surgery related to COVID  
☐ Yes, chose to avoid open surgery related to COVID

**Surgical intent:** Curative | Palliative

**Environment in which patient cared for:**

**Operative theatre** – Designated COVID theatre | Designated non-COVID theatre | No designation for theatre

**ITU** – Designated COVID treatment area | Designated non-COVID treatment area | No designation for this area

**Postop ward** – Designated COVID ward | Designated non-COVID ward | No designation for this ward

**Did any change to treatment occur due to the COVID-19 pandemic (operated patients)?**

No change to care, no neoadjuvant Rx | No change – neoadjuvant equivalent to pre-COVID | Delay to definitive Rx | Expedited definitive surgery | Change in choice of operation | Op in alt. hospital | IR before surgery, not typically indicated | Neoadj treatment, not typically indicated | No Neoadj treatment, typically indicated | Neoadj treatment longer than typical | Neoadj treatment shorter than typical | Adj treatment, not typically indicated | No Adj treatment, typically indicated | Other: \_\_\_\_\_

Neoadj Rx: Radiotherapy | Chemotherapy | Other: \_\_\_\_\_

Details of neoadj Rx: \_\_\_\_\_

**COVID-19 post-operatively** (30 days): Yes – lab test | Yes – CT thorax | Yes – clinical only | No

If yes: Inpatient | Required Admission | Community

**Mortality:** Died on table | d0-7 | d8-30 |

Alive still in hosp 30d | transferred | discharged to rehab | discharged home

**Re-operation:** Yes | No

**Post-op intensive care:** No | planned from theatre | unplanned from theatre | unplanned from ward  
(If no/unplanned from ward): **Would post-operative ICU bed have been planned pre-COVID-19 era?** Yes, not available ~ COVID | Yes, not available (other) | No

**Total length of hospital stay:** \_\_\_\_\_ days

**Did surgeons contract COVID-19** (30-days): Yes | No

**If a cancer operation WAS performed:**

**Complications:**

- |   |   |
|---|---|
| <input type="checkbox"/> Acute kidney injury        | <input type="checkbox"/> Pneumonia            |
| <input type="checkbox"/> ARDS                       | <input type="checkbox"/> Respiratory failure  |
| <input type="checkbox"/> Anastomotic leak           | <input type="checkbox"/> Sepsis               |
| <input type="checkbox"/> Blood transfusion          | <input type="checkbox"/> Septic shock         |
| <input type="checkbox"/> Cardiac arrest             | <input type="checkbox"/> Stroke/TIA           |
| <input type="checkbox"/> Coma >24h                  | <input type="checkbox"/> SSI superficial/deep |
| <input type="checkbox"/> Deep Vein Thrombosis       | <input type="checkbox"/> SSI organ space      |
| <input type="checkbox"/> Graft/prosthesis/flap fail | <input type="checkbox"/> UTI                  |
| <input type="checkbox"/> Myocardial infraction      | <input type="checkbox"/> Wound dehiscence     |

**If NO operation was performed (by 3 months from study entry)**

Is there **still** a plan for curative surgery? Yes | No

**Why** was no operation performed in the 3 months?

**If still plan for surgery:**

- ☐ Patient choice to avoid surgery during pandemic  
☐ MDT decision to delay surgery due to risk to patient  
☐ Ongoing neoadjuvant treatment  
☐ No bed / intensive care space / theatre space  
☐ Change of recommendations in society guidelines related to COVID-19

**If no ongoing plan for surgery:**

- ☐ Patient choice to avoid surgery during pandemic  
☐ MDT decision to delay surgery due to risk to patient  
☐ Disease progression, surgery no longer indicated  
☐ Change in clinical status unrelated to cancer e.g. MI  
☐ Died awaiting surgery  
☐ Changes in society guidelines due to COVID-19  
☐ Other: \_\_\_\_\_

Has the cancer been **re-staged**? No | Yes, If so, date \_\_\_\_/\_\_\_\_/\_\_\_\_

**NCCN classification:**

Resectable | Borderline resectable (vein involvement) | Borderline resectable (arterial involvement) | Locally advanced | Metastatic

**Did any change to treatment occur due to the COVID-19 pandemic (non-operated patients)?**

No change to care – delayed/cancelled other reason | Operation cancelled because of COVID-19 | Operation delayed because of COVID-19 | Change in Rx strategy due to COVID-19 | IR before/instead of surgery, not typically indicated | Neoadj treatment, not typically indicated | No Neoadj treatment, typically indicated | Neoadj treatment longer than typical | Neoadj treatment shorter than typical | Less access to staging procedures | Less access to staging investigations | Other: \_\_\_\_\_

Neoadj Rx: Radiotherapy | Chemotherapy | Other: \_\_\_\_\_

Details of neoadj Rx: \_\_\_\_\_


**Unplanned admission with cholangitis**

- ☐ Yes ☐ No



## Case Report Form CovidSurg

NB: Complete this CRF only for patients that are eligible for **both** studies (operated cancer patients with COVID-19 infection)



Patient REDCap ID: \_\_\_\_\_

### COVID-19 Patient Information

#### BCG/Tuberculosis(TB) status

Vaccine - <15yrs ago | Vaccine – 15>yrs ago | TB diagnosis <15yrs ago | TB diagnosis >15yrs ago | Close contact with individual with known TB <15yrs ago | Close contact with individual with known TB >15yrs ago | No previous BCG vaccine/TB exposure | BCG vaccination/TB exposure unknown

#### Findings at admission:

- |   |  |
|---|--|
| <input type="checkbox"/> Abdominal pain | <input type="checkbox"/> Haemoptysis     |
| <input type="checkbox"/> Dyspnoea       | <input type="checkbox"/> Myalgia         |
| <input type="checkbox"/> Cough          | <input type="checkbox"/> Nausea/vomiting |
| <input type="checkbox"/> Diarrhoea      | <input type="checkbox"/> Sputum          |
| <input type="checkbox"/> Fatigue        | <input type="checkbox"/> Other: _____    |
| <input type="checkbox"/> Fever >38C     |  |

### COVID-19 Preoperative Investigations

#### Last available data from before surgery:

Resp rate: \_\_\_\_\_ rpm Heart rate: \_\_\_\_\_ bpm  
Systolic BP: \_\_\_\_\_ mmHg Diastolic BP: \_\_\_\_\_ mmHg

#### Tests performed to investigate SARS-CoV-2 status:

	CT (neg)	CT (pos)	Swab (neg)	Swab (pos)
4-7 days prior surgery	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
1-3 days prior surgery	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Day of surgery (preop)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
After surgery (during index admission)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
After discharge from index admission (within 30 days of surgery)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

#### How was SARS-CoV-2 confirmed?

- ☐ Positive swab – result received before surgery  
☐ Positive swab – result received after surgery  
☐ CT scan of chest confirming COVID-19 – before surgery  
☐ CT scan of chest confirming COVID-19 – after surgery  
☐ Clinical diagnosis/chest X-Ray – before surgery  
☐ Clinical diagnosis/chest X-Ray – after surgery

#### Pre-op investigations:

Haemoglobin: \_\_\_\_\_ g/L WCC: \_\_\_\_\_ 10<sup>9</sup>/L CRP: \_\_\_\_\_ mg/L

#### Pre-op x-ray:

- ☐ Not performed  
☐ Yes- normal  
☐ Yes- abnormal

#### Pre-op chest CT:

- ☐ Not performed  
☐ Yes- normal  
☐ Yes- consolidation  
☐ Yes- ground glass opacity  
☐ Yes- pulmonary infiltration  
☐ Yes- other abnormality

### Management

#### Time from admission to operation (pre-op delay)

<6 hrs | 6-23 hrs | 24-47 hrs | 48-71 hrs | 72+hrs

### COVID-19 Treatment

Did patient receive NSAIDs? No | Yes before admission  
 | After admission | Both

#### Patient received during index admission?

- |   |   |
|---|---|
| <input type="checkbox"/> Antibiotics        | <input type="checkbox"/> IV Immunoglobulins |
| <input type="checkbox"/> Antivirals         | <input type="checkbox"/> Interferon         |
| <input type="checkbox"/> Quinine/derivative | <input type="checkbox"/> IL-6 blocker       |
| <input type="checkbox"/> Corticosteroids    |   |

Antiviral (name & dose): \_\_\_\_\_

Corticosteroid (name & dose): \_\_\_\_\_

#### Renal dialysis during index admission?

No | Yes but not at 30 days after surgery  
 Yes and ongoing dialysis at 30 days after surgery

#### Pre-op respiratory support:

- |                                       |   |
|---------------------------------------|---|
| <input type="checkbox"/> None         | <input type="checkbox"/> Non-invasive ventilation |
| <input type="checkbox"/> Low-flow O2  | <input type="checkbox"/> Invasive vent            |
| <input type="checkbox"/> High-flow O2 | <input type="checkbox"/> ECMO                     |

#### Post-op respiratory support:

- |                                       |   |
|---------------------------------------|---|
| <input type="checkbox"/> None         | <input type="checkbox"/> Non-invasive ventilation |
| <input type="checkbox"/> Low-flow O2  | <input type="checkbox"/> Invasive vent            |
| <input type="checkbox"/> High-flow O2 | <input type="checkbox"/> ECMO                     |

#### Duration of post-op mechanical ventilation:

1-23h | 24-47h | 48-71h | 72-167h | 168h+